

Effectiveness of woodland birds as taxonomic surrogates in conservation planning for biodiversity on farms

Karen Ikin¹, Ding Li Yong¹ and David Lindenmayer¹

¹Fenner School of Environment and Society, The Australian National University, Frank Fenner Building 141, Linnaeus Way, Acton ACT 2601, Australia

ARC Centre of Excellence for Environmental Decisions, The Australian National University, Frank Fenner Building 141, Linnaeus Way, Acton ACT 2601, Australia

National Environmental Science Program Threatened Species Hub, The Australian National University, Frank Fenner Building 141, Linnaeus Way, Acton ACT 2601, Australia

Correspondence: karen.ikin@anu.edu.au Fenner School of Environment and Society, The Australian National University, Linnaeus Way, Acton ACT 2601, Australia. +61 2 6125 1494

Article type: Research paper

Abstract

Woodland birds are a commonly used taxonomic surrogate for other species groups in agricultural landscapes as they are relatively diverse, easily-studied, and charismatic. Yet, other taxa can respond to native vegetation on farms differently to woodland birds, challenging the present focus on birds in agri-environmental schemes. We aimed to assess the effectiveness of woodland birds as taxonomic surrogates for biodiversity in conservation planning on farms, in comparison with reptiles and arboreal marsupials. We used a complementarity-based approach to select patches of remnant and restored vegetation that supported *a priori* representation targets of species occurrences. We found that the spatial locations of vegetation patches selected to meet representation targets for woodland birds were 24% - 69% different from the locations of patches selected for other taxa. The vegetation patches selected to meet representation targets for woodland birds failed to incidentally meet representation targets for other taxa, although targets for a subset of threatened woodland birds were exceeded. Conservation planning for woodland birds, however, led to higher incidental representation of the other taxa, compared with conservation planning for reptiles and arboreal marsupials. This indicates that woodland birds are a more effective taxonomic surrogate for biodiversity on farms compared to reptiles and arboreal marsupials. If the conservation goal is to conserve a broad array of biodiversity on farms, then the focus on woodland birds in agri-environmental schemes is justified. However, if the conservation of particular species or taxonomic groups is a priority, then conservation plans explicitly targeting these species or groups are required.

Keywords: Agricultural landscapes, Arboreal marsupials, Incidental representation, Marxan, Reptiles, Woodland birds

24 **Highlights**

- 25 • We compared woodland birds, reptiles, arboreal marsupials as taxonomic surrogates
- 26 • Conservation planning for any one taxon failed to incidentally represent other taxa
- 27 • Yet, woodland birds were more effective taxonomic surrogates than other taxa
- 28 • Threatened bird species were represented by woodland bird conservation plans
- 29 • Focus on woodland birds can conserve a broad array of biodiversity on farms

30

1. Introduction

A core challenge for conservation science is the lack of complete information on biodiversity, that is, a comprehensive inventory of all species of all groups in a given area (Williams and Gaston 1994). This challenge is difficult to address directly, given insufficient resources to survey the myriad of species in ecosystems, as well as the spatial and temporal complexity of ecosystem processes. Instead, surrogates for biodiversity are used, for instance environmental attributes or taxonomic groups, that attempt to represent the full assemblages of species to some degree (Howard et al. 1998; Andelman and Fagan 2000; Margules and Pressey 2000; Sarkar et al. 2006; Rodrigues and Brooks 2007).

Birds are the most commonly used taxonomic surrogate in terrestrial ecosystems (Eglington et al. 2012; Larsen et al. 2012; Westgate et al. 2014). They are a well-studied taxon, being highly detectable, easily identifiable, and inexpensive to survey compared with other vertebrate and invertebrate taxa. Their relatively high levels of species diversity, breadth of functional attributes, and heterogeneous distributions also contribute to their effectiveness in improving the efficiency of conservation planning and management (Lewandowski et al. 2010). Further, birds are a charismatic taxon garnering high public appeal, which makes them an ideal flagship group for conservation actions (Veríssimo et al. 2009).

In agricultural landscapes, birds are often the target group for agri-environmental initiatives (Guerrero et al. 2012), including restoration plantings and the protection of remnant vegetation. In Australia, most restoration initiatives aimed at improving biodiversity conservation (e.g. Lindenmayer et al. 2013) have focused on woodland birds. Woodland birds are defined here as species that occur in temperate woodland, not excluding species that also occur in grassland (Silcocks et al. 2005). There is a vast literature on woodland birds, exploring the importance of different vegetation attributes at patch and landscape scales (e.g. Watson et al. 2003; Radford et al. 2005; Barrett et al. 2008; Haslem and Bennett 2008; Bowen et al. 2009; Hanspach et al. 2011; Ikin et al. 2014), and in conservation planning (Thomson et al. 2009; Ikin et al. 2016). Findings from these studies contribute to the evidence-base for conserving a broad array of biodiversity on farms. However, other research shows that other groups of vertebrate taxa that are more difficult to survey, for example mammals and reptiles, can respond differently to vegetation composition and structure compared to woodland birds

(Cunningham et al. 2007; Jellinek et al. 2014; Michael et al. 2014; Yong et al. 2016). Such a discrepancy in responses to the landscape calls into question whether woodland birds are as good taxonomic surrogates for biodiversity on farms as they are supposed.

Our study aimed to assess the effectiveness of woodland birds as taxonomic surrogates in conservation planning for biodiversity on farms, in comparison with reptiles and arboreal marsupials. Agricultural landscapes, despite their highly modified state, can support high numbers of species (Yong et al. 2016), and systematic survey data on multiple taxonomic groups are rare (underscoring the necessity of using taxonomic surrogate approaches). We took advantage of the South West Slopes Restoration Study (Cunningham et al. 2007; Lindenmayer et al. 2016), which gathers detailed multi-taxon data across an extensive agricultural region of southeastern Australia. Using a complementarity-based approach, for each taxonomic group we identified patches of restored and remnant vegetation that together met *a priori* representation targets of species occurrence in the landscape. We asked:

1. Are the vegetation patches selected to meet representation targets for one taxon the same as vegetation patches selected for other taxa?
2. Which taxon achieved the best incidental representation of other taxa?

Given the relatively high species diversity and functional diversity of birds in the landscape, we predicted that a large number of restored and remnant vegetation patches would be needed to meet niche requirements (Moritz et al. 2001). Consequently, we expected that spatial concordance between these patches and those selected to meet representation targets for other taxa would be high, and therefore that conservation planning for woodland birds would be effective at incidentally representing other taxa – thus indicating that woodland birds are effective surrogates. However, we did not expect that the degree of spatial concordance or incidental representation would be identical between taxa (Lentini and Wintle 2015). For instance, we predicted that conservation planning for all woodland birds in our agricultural landscape would be: (i) less effective than conservation planning targeted at a subset of threatened woodland birds (Beger et al. 2003), and (ii) more effective at representing arboreal marsupials, which may use the landscape at similar scales to birds and thus may have similar ecological requirements, compared to reptiles, which may use the landscape at smaller scales (Yong et al. 2016).

2. Methods

2.1 Study design

We conducted our study in a 150 km x 120 km area of the wheat-sheep belt of southeastern Australia, in the South West Slopes bioregion of New South Wales (Fig. A1 in the supplementary material). Farms within this region typically have between 3% and 35% native vegetation cover, including remnant temperate box-gum *Eucalyptus* woodland, natural and coppiced regrowth, and restoration plantings (Cunningham et al. 2014). For this investigation, we focused on 189 patches of native vegetation (68 remnant woodland, 61 regrowth woodland, and 60 plantings), which together covered 1437 ha across 43 farms (Table A1 in the supplementary material).

We collected two years of occurrence data for each of our taxonomic groups along a permanent 200 m transect established in each patch. Bird surveys were conducted in spring 2008 and 2011, with each transect visited twice in any given year between sunrise and mid-morning. Each visit involved five-minute point counts at the 0 m, 100 m and 200 m transect points. All birds seen or heard within 50 m of the point, but excluding those flying overhead, were recorded as present. Reptile surveys were conducted in spring 2008 and winter 2011, with each transect visited once between mid-morning and mid-afternoon. Each visit involved a twenty-minute active search of leaf litter, grass tussocks, coarse woody debris, surface rocks, and exfoliating bark, between the 0 m and 200 m transect points. All reptiles seen within 50 m were recorded as present. Visits also involved inspecting arrays of artificial refuges (four wooden railway sleepers, four terracotta roof tiles, and one double stack of 1-m² corrugated steel sheet) placed at the 0 m and 100 m transect points. Arboreal marsupial surveys were conducted in autumn 2008 and winter 2011, with each transect visited once between sunset and midnight. Each visit involved a twenty-minute spotlight survey between the 100 m and 200 m transect points, walking at an average speed of 3 km/h. All species seen or heard were recorded as present.

2.2 Data analysis

We restricted our analysis to species recorded at least twice over the two survey years (Table 1; Table A2 in the supplementary material). This enabled us to exclude vagrant species. This gave 72 species of woodland birds (Silcocks et al. 2005); a subset of 10 species of listed birds (woodland birds listed as threatened in New South Wales in 2016 under the *Threatened Species Conservation Act 1995*; hereafter referred to as a separate taxon for simplicity); three species of arboreal marsupials; and 22 species of reptiles.

For each taxonomic group, our objective was to find complementary sets of patches that met *a priori* representation targets of species occurrences while minimizing the combined area (ha) of the patch set, irrespective of spatial configuration (note that this objective of minimizing the area of vegetation needed to meet representation targets is not intended to identify “unnecessary” vegetation patches, but instead constrain the analyses to best compare surrogate efficacy). To do this, we used Marxan, a decision-support software program that uses a simulated annealing algorithm to solve the minimum set problem (Ball et al. 2009). We created a conservation feature representing patch occurrence of each species in each survey year (two features per species, e.g. for woodland birds we created 144 conservation features in total), following Ikin et al. (2016) and Runge et al. (2016). We set representation targets of 25%, 50%, and 75% occurrence of species in every year (equivalent to 25%, 50%, and 75% of patches where each species occurred). For every combination of taxon and representation target (12 in total), we performed 100 Marxan runs to identify the best patch set. The best patch set was defined as selected patches of vegetation that represented the target of species occurrences in the landscape over the two study years (e.g. 25% representation of woodland bird species occurrences, while ignoring the occurrences of arboreal marsupials and reptiles) for the least combined area. To confirm that patch selection for woodland birds was not sensitive to the subset of listed birds, we re-ran the analyses for woodland birds excluding listed species.

To answer our first question (*Are the vegetation patches selected to meet representation targets for one taxon the same as vegetation patches selected for other taxa?*), we assessed the spatial concordance between the best patch sets for each taxon and representation target. To do this, we calculated Bray-Curtis dissimilarity (adjusted for presence-absence data) between each pair of best

patch sets, with low dissimilarity indicating that the spatial locations of the selected patches were similar.

To answer our second question (*Which taxon achieved the best incidental representation of other taxa?*), we assessed how well the best patch sets selected for one taxon represented the occurrences of species in each of the other three taxa. To do this, we calculated the average minimum percent occurrence of each species per taxon that was met over the study period under each best patch set. Incidental representation is a direct measure of surrogate efficacy (Grantham et al. 2010) – the higher the incidental representation of other taxa a particular taxon achieves, the more effective that taxon is as a taxonomic surrogate.

3. Results

Woodland birds were the most species-diverse taxon of the three taxa we studied, every study patch supported at least one woodland bird species, and each species occurred in a median of 10.25 patches (Table 1). In comparison, arboreal marsupials were the least species-diverse taxon, only 51% of patches supported at least one arboreal marsupial species, and each species occurred in a median of 38.00 patches. Consistently across representation targets (25%, 50%, and 75% species occurrences in 2008 and 2011), we found the combined area of the vegetation patches that represented target occurrences of species in the landscape for the least combined area (i.e. the best patch sets) was largest for woodland birds and smallest for arboreal marsupials, although the relative difference in area decreased as representation targets increased from 25% to 75% of species occurrences (Table 1).

We found considerable difference between the spatial locations of the patches in the best patch sets for each taxon and representation target (Fig. 1). For example, the locations of patches in the best patch set to achieve the 25% representation target for arboreal marsupials was up to 76% different from the locations of patches in the best patch sets that met this representation target for other taxa. Even between all woodland birds versus the subset of listed woodland birds, there was up to 55% difference in the locations of patches in the best patch sets selected to meet the same representation target. Similarity between the locations of the patches in the best patch sets was highest between woodland birds and reptiles (as low as 24% difference for the 75% representation target).

In general, we found that the best patch sets selected for one taxon failed to meet representation targets for other taxa (Fig. 2). The best patch sets for woodland birds, as an exception, exceeded targets for the occurrences of the subset of listed woodland birds (Fig. 2a). These best patch sets also came close to meeting target occurrences of the other taxa. For instance, the best patch set to meet the 75% representation target also represented 73% of arboreal marsupials and 69% of reptiles. Listed birds were not driving these patterns as results were similar when this subset of species was removed from the woodland bird taxon (Fig. A2 in the supplementary material). The best patch sets for listed birds, in contrast, did not meet representation targets for other woodland birds, nor representation targets for the other taxa (Fig. 2b). The best patch sets for arboreal marsupials were the worst for representing the occurrences of other taxa; for instance, the best patch set selected to achieve the 75% representation target for arboreal marsupials represented only 27% of woodland birds, 37% of listed birds and 25% of reptiles. (Fig. 2c).

4. Discussion

Woodland birds are a commonly used taxonomic surrogate for other species groups in agricultural landscapes (Eglington et al. 2012; Larsen et al. 2012), but how do they compare with arboreal marsupials and reptiles in conservation planning for biodiversity on farms? We found that the spatial locations of the best sets of vegetation patches selected to meet representation targets for woodland birds were between 24% and 69% different from the locations of the best patch sets selected for other taxa. The locations of the best patch sets selected for reptiles showed a similar amount of spatial concordance to woodland birds, but those selected for arboreal marsupials were between 46% and 76% different from other best patch sets. We found that the best patch sets selected to meet representation targets for woodland birds failed to incidentally meet representation targets for other taxa, although targets for the subset of threatened woodland birds were exceeded. Conservation planning for woodland birds, however, led to higher incidental representation of the other taxa (up to 73% representation under the 75% representation target), compared with conservation planning for arboreal marsupials (up to 27%) and reptiles (up to 62%).

The high species diversity of woodland birds, coupled with the relatively low median number of patches occupied by each species, contributed to their effectiveness as taxonomic surrogates in our study system. In contrast, the species-poor but widely-distributed arboreal marsupial taxon was the least effective taxonomic surrogate. Previous studies have found similar relationships between surrogate efficacy, species diversity and spatial distribution. For example, Beger et al. (2003) demonstrated that marine reserves developed for fish species with heterogeneous distributions were better at representing coral species with homogenous distributions, than vice versa. Similarly, Moritz et al. (2001) found the high diversity and narrow distributions of rainforest invertebrates made them better surrogates compared with less diverse, more broadly distributed taxa. While many studies have assessed vertebrates, those that test the effectiveness of invertebrate taxa as surrogates remain fairly limited (Sauberer et al. 2004), in spite of the ecological importance of many insect groups. We suggest that future research consider how conservation plans for vertebrate taxa, such as birds, represent invertebrates of explicit importance to farm production (e.g. bees), and also if conservation planning for biodiversity on farms can be improved through incorporating non-vertebrate groups.

The broader range of vegetation niches occupied by woodland birds also made them better taxonomic surrogates than arboreal marsupials and reptiles. For example, among the woodland bird taxon were species that foraged in leaf litter (e.g. speckled warbler, *Chthonicola sagittata*), under decorticating bark (e.g. crested shrike-tit, *Falcunculus frontatus*), and in the tree canopy (e.g. striated thornbill, *Acanthiza lineata*); species that fed on invertebrates (e.g. grey fantail, *Rhipidura albiscapa*), nectar (e.g. little lorikeet, *Glossopsitta pusilla*), and seeds (e.g. crested pigeon, *Ocyphaps lophotes*); and species that nested in the understory (e.g. superb fairy wren, *Malurus cyaneus*) and tree cavities (e.g. crimson rosella, *Platycercus elegans*). Thus, the ecological requirements of woodland birds overlapped with those of the arboreal marsupials and reptiles, but the reverse was not true; the arboreal marsupials, in particular, had very low niche diversity (perhaps explaining the high spatial difference between the best patch sets selected for this taxon compared with the other taxa). Our results support previous studies that have found that taxa with similar dependencies on their environment make better surrogates for each other compared with taxa that have different dependencies (Howard et al. 1998; Mortelliti et al. 2008; Heino et al. 2009).

Conservation planning for woodland birds was effective at representing the subset of threatened woodland birds, with representation targets exceeded even when the threatened species were not explicitly considered in the plans. This result was unexpected, as threatened species generally have more restricted distributions, making them more likely to be unrepresented in conservation landscapes (Moore et al. 2003; Grantham et al. 2010). Myšák and Horsák (2014), for example, found that the species richness of red-listed cryptogams and snails were poor surrogates for the species richness of all cryptogams and snails and vice versa. However, consistent with their study, we found that vegetation patches selected to represent threatened woodland birds did not meet targets for other woodland birds, nor arboreal marsupials and reptiles.

We incorporated two years of species occurrence data in our analyses, including from severe drought (2008) and post-drought recovery (2011). This approach accounts for variance in species distributions over time (Ikin et al. 2016; Runge et al. 2016), and thus may improve the robustness of conservation plans to stochastic disturbances (Lourival et al. 2011; Van Teeffelen et al. 2012). However, by only considering species *representation* across the landscape, it is difficult to determine the efficacy of each taxon as surrogates for species *persistence*. It is possible that focusing conservation planning on the population viability of at-risk species, e.g. the group of listed woodland birds, will lead to improved conservation outcomes for other taxa (Williams and Araújo 2000; Nicholson et al. 2013). Thus, we acknowledge it is possible that assessing the effectiveness of taxonomic surrogates using incidental persistence instead of incidental representation would give a different conclusion as to which taxa was the best surrogate. Future research should consider this question, perhaps using new methods that incorporate both representation and persistence in conservation plans (e.g. Bode et al. 2016).

Our study demonstrates the fundamental trade-offs inherent in single-taxon conservation planning, and taxonomic surrogate approaches (Andelman and Fagan 2000; Wiens et al. 2008). Representation targets for individual taxa were met only through taxon-specific conservation plans, but these plans failed to represent broader farmland biodiversity. Woodland birds proved the best taxonomic surrogates (despite failing to meet targets for arboreal marsupials and reptiles) but the sets of vegetation patches selected to meet representation targets for this taxon were the most spatially

extensive. Given that farmland prioritized for biodiversity conservation may compromise production opportunities, spatially extensive conservation plans in these landscapes may have serious economic consequences and may not be feasible or cost-effective to implement or manage (House et al. 2008). In comparison, representation targets for threatened woodland birds could be met with less than 50% of the vegetation area required, but few species from other taxa were also fully represented. It is also important to note that approximately 85% of temperate woodland has been cleared from our study region (Benson 2008), and all remaining vegetation contributes to conservation outcomes (Cunningham et al. 2014). Incidental representation could be improved by incorporating additional species or taxa into the conservation plans (Moore et al. 2003; Larsen et al. 2012), but this approach may increase farmland area prioritized for conservation and thus also increase opportunity costs associated with lost production. These conundrums are not easy to resolve, but require *a priori* value judgements of which aspects of biodiversity on farms should be conserved and what management considerations also should be taken into account.

In conclusion, our study shows that the diverse, easily-studied, and charismatic woodland bird taxon is a more effective taxonomic surrogate than other major farmland vertebrate taxa in this landscape. The present focus on woodland birds in agri-environmental schemes (Guerrero et al. 2012) is thus justified if the conservation goal is to conserve a broad array of biodiversity on farms. However, if particular species or taxonomic groups are considered a conservation priority, then conservation plans explicitly targeting these species or groups will be required to meet conservation goals.

Acknowledgements

We thank the Australian Research Council, the Murray and Riverina Local Land Services, and the Caring for Our Country Program for funding for this project. DLY thanks the Leslie Foundation for support. We thank the large number of people within the Conservation and Landscape Ecology Group who helped collect the field data or provided comments on the paper.

References

- Andelman, S.J., Fagan, W.F., 2000. Umbrellas and flagships: Efficient conservation surrogates or expensive mistakes? *Proceedings of the National Academy of Sciences* 97, 5954-5959.
- Barrett, G., Freudenberger, D., Drew, A., Stol, J., Nicholls, A.O., Cawsey, E.M., 2008. Colonisation of native tree and shrub plantings by woodland birds in an agricultural landscape. *Wildlife Research* 35, 19-32.
- Beger, M., Jones, G.P., Munday, P.L., 2003. Conservation of coral reef biodiversity: a comparison of reserve selection procedures for corals and fishes. *Biological Conservation* 111, 53-62.
- Benson, J., 2008. New South Wales Vegetation Classification and Assessment: Part 2 Plant communities of the NSW South-western Slopes Bioregion and update of NSW Western Plains plant communities, Version 2 of the NSWVCA database. *Cunninghamia* 104, 599-673.
- Bode, M., Williamson, D.H., Weeks, R., Jones, G.P., Almany, G.R., Harrison, H.B., Hopf, J.K., Pressey, R.L., 2016. Planning Marine Reserve Networks for Both Feature Representation and Demographic Persistence Using Connectivity Patterns. *PLoS ONE* 11, e0154272.
- Bowen, M.E., McAlpine, C.A., Seabrook, L.M., House, A.P.N., Smith, G.C., 2009. The age and amount of regrowth forest in fragmented brigalow landscapes are both important for woodland dependent birds. *Biological Conservation* 142, 3051-3059.
- Cunningham, R., Lindenmayer, D.B., Michael, D., Crane, M., Barton, P.S., Gibbons, P., Okada, S., Ikin, K., Stein, J.A.R., 2014. The law of diminishing returns: woodland birds respond to native vegetation cover at multiple spatial and temporal scales *Diversity and Distributions* 20, 59–71.
- Cunningham, R.B., Lindenmayer, D.B., Crane, M., Michael, D., MacGregor, C., 2007. Reptile and arboreal marsupial response to replanted vegetation in agricultural landscapes. *Ecological Applications* 17, 609-619.

302 Eglinton, S.M., Noble, D.G., Fuller, R.J., 2012. A meta-analysis of spatial relationships in species
 303 richness across taxa: Birds as indicators of wider biodiversity in temperate regions. *Journal for Nature*
 304 *Conservation* 20, 301-309.

305 Grantham, H.S., Pressey, R.L., Wells, J.A., Beattie, A.J., 2010. Effectiveness of Biodiversity
 306 Surrogates for Conservation Planning: Different Measures of Effectiveness Generate a Kaleidoscope
 307 of Variation. *PLoS ONE* 5, e11430.

308 Guerrero, I., Morales, M.B., Oñate, J.J., Geiger, F., Berendse, F., Snoo, G.d., Eggers, S., Pärt, T.,
 309 Bengtsson, J., Clement, L.W., Weisser, W.W., Olszewski, A., Ceryngier, P., Hawro, V., Liira, J.,
 310 Aavik, T., Fischer, C., Flohre, A., Thies, C., Tschardtke, T., 2012. Response of ground-nesting
 311 farmland birds to agricultural intensification across Europe: Landscape and field level management
 312 factors. *Biological Conservation* 152, 74-80.

313 Hanspach, J., Fischer, J., Stott, J., Stagoll, K., 2011. Conservation management of eastern Australian
 314 farmland birds in relation to landscape gradients. *Journal of Applied Ecology* 48, 523-531.

315 Haslem, A., Bennett, A.F., 2008. Birds in agricultural mosaics: the influence of landscape pattern and
 316 countryside heterogeneity. *Ecological Applications* 18, 185-196.

317 Heino, J., Tolonen, K.T., Kotanen, J., Paasivirta, L., 2009. Indicator groups and congruence of
 318 assemblage similarity, species richness and environmental relationships in littoral macroinvertebrates.
 319 *Biodiversity and Conservation* 18, 3085-3098.

320 House, A.P.N., MacLeod, N.D., Cullen, B., Whitbread, A.M., Brown, S.D., McIvor, J.G., 2008.
 321 Integrating production and natural resource management on mixed farms in eastern Australia: The
 322 cost of conservation in agricultural landscapes. *Agriculture, Ecosystems & Environment* 127, 153-
 323 165.

324 Howard, P.C., Viskanic, P., Davenport, T.R.B., Kigenyi, F.W., Baltzer, M., Dickinson, C.J., Lwanga,
 325 J.S., Matthews, R.A., Balmford, A., 1998. Complementarity and the use of indicator groups for
 326 reserve selection in Uganda. *Nature* 394, 472-475.

327 Ikin, K., Barton, P.S., Stirnemann, I.A., Stein, J.R., Michael, D., Crane, M., Okada, S., Lindenmayer,
 328 D.B., 2014. Multi-Scale Associations between Vegetation Cover and Woodland Bird Communities
 329 across a Large Agricultural Region. *PLoS ONE* 9, e97029.

330 Ikin, K., Tulloch, A., Gibbons, P., Ansell, D., Seddon, J., Lindenmayer, D., 2016. Evaluating
 331 complementary networks of restoration plantings for landscape-scale occurrence of temporally-
 332 dynamic species. *Conservation Biology* 30, 1027–1037.

333 Jellinek, S., Parris, K.M., McCarthy, M.A., Wintle, B.A., Driscoll, D.A., 2014. Reptiles in restored
 334 agricultural landscapes: the value of linear strips, patches and habitat condition. *Animal Conservation*
 335 17, 544-554.

336 Larsen, F.W., Bladt, J., Balmford, A., Rahbek, C., 2012. Birds as biodiversity surrogates: will
 337 supplementing birds with other taxa improve effectiveness? *Journal of Applied Ecology* 49, 349-356.

338 Lentini, P.E., Wintle, B.A., 2015. Spatial conservation priorities are highly sensitive to choice of
 339 biodiversity surrogates and species distribution model type. *Ecography* 38, 1101-1111.

340 Lewandowski, A.S., Noss, R.F., Parsons, D.R., 2010. The Effectiveness of Surrogate Taxa for the
 341 Representation of Biodiversity. *Conservation Biology* 24, 1367-1377.

342 Lindenmayer, D., Willinck, E., Crane, M., Michael, D., Okada, S., Cumming, C., Durant, K.,
 343 Frankenberg, J., 2013. Murray Catchment habitat restoration: Lessons from landscape-level research
 344 and monitoring. *Ecological Management & Restoration* 14, 80-92.

345 Lindenmayer, D.B., Lane, P.W., Barton, P.S., Crane, M., Ikin, K., Michael, D., Okada, S., 2016.
 346 Long-term bird colonization and turnover in restored woodlands. *Biodiversity and Conservation* 25,
 347 1587-1603.

348 Lourival, R., Drechsler, M., Watts, M.E., Game, E.T., Possingham, H.P., 2011. Planning for reserve
 349 adequacy in dynamic landscapes; maximizing future representation of vegetation communities under
 350 flood disturbance in the Pantanal wetland. *Diversity and Distributions* 17, 297-310.

351 Margules, C.R., Pressey, R.L., 2000. Systematic conservation planning. *Nature* 405, 243-253.

352 Michael, D.R., Wood, J.T., Crane, M., Montague-Drake, R., Lindenmayer, D.B., 2014. How effective
 353 are agri-environment schemes for protecting and improving herpetofaunal diversity in Australian
 354 endangered woodland ecosystems? *Journal of Applied Ecology*, DOI: 10.1111/1365-2664.12215.

355 Moore, J.L., Balmford, A., Brooks, T., Burgess, N.D., Hansen, L.A., Rahbek, C., Williams, P.H.,
 356 2003. Performance of Sub-Saharan Vertebrates as Indicator Groups for Identifying Priority Areas for
 357 Conservation

358 La Efectividad de los Vertebrados del Sub-Sahara como Grupos Indicadores para Identificar Áreas
 359 Prioritarias para la Conservación. *Conservation Biology* 17, 207-218.

360 Moritz, C., Richardson, K.S., Ferrier, S., Monteith, G.B., Stanisci, J., Williams, S.E., Whiffin, T.,
 361 2001. Biogeographical Concordance and Efficiency of Taxon Indicators for Establishing
 362 Conservation Priority in a Tropical Rainforest Biota. *Proceedings: Biological Sciences* 268, 1875-
 363 1881.

364 Mortelliti, A., Santulli Sanzo, G., Boitani, L., 2008. Species' surrogacy for conservation planning:
 365 caveats from comparing the response of three arboreal rodents to habitat loss and fragmentation.
 366 *Biodiversity and Conservation* 18, 1131-1145.

367 Myšák, J., Horsák, M., 2014. Biodiversity surrogate effectiveness in two habitat types of contrasting
 368 gradient complexity. *Biodiversity and Conservation* 23, 1133-1156.

369 Nicholson, E., Lindenmayer, D.B., Frank, K., Possingham, H.P., 2013. Testing the focal species
 370 approach to making conservation decisions for species persistence. *Diversity and Distributions* 19,
 371 530-540.

372 Radford, J.Q., Bennett, A.F., Cheers, G.J., 2005. Landscape-level thresholds of habitat cover for
373 woodland-dependent birds. *Biological Conservation* 124, 317-337.

374 Rodrigues, A.S.L., Brooks, T.M., 2007. Shortcuts for Biodiversity Conservation Planning: The
375 Effectiveness of Surrogates. *Annual Review of Ecology, Evolution, and Systematics* 38, 713-737.

376 Runge, C.A., Tulloch, A., Possingham, H., Tulloch, V., Fuller, R.A., 2016. Incorporating dynamic
377 distributions into spatial prioritization. *Diversity & Distributions* 22, 332–343

378 Sarkar, S., Pressey, R.L., Faith, D.P., Margules, C.R., Fuller, T., Stoms, D.M., Moffett, A., Wilson,
379 K.A., Williams, K.J., Williams, P.H., Andelman, S., 2006. Biodiversity Conservation Planning Tools:
380 Present Status and Challenges for the Future. *Annual Review of Environment and Resources* 31, 123-
381 159.

382 Sauberer, N., Zulka, K.P., Abensperg-Traun, M., Berg, H.-M., Bieringer, G., Milasowszky, N.,
383 Moser, D., Plutzer, C., Pollheimer, M., Storch, C., Tröstl, R., Zechmeister, H., Grabherr, G., 2004.
384 Surrogate taxa for biodiversity in agricultural landscapes of eastern Austria. *Biological Conservation*
385 117, 181-190.

386 Silcocks, A., Tzaros, C., Weston, M., Olsen, P., 2005. An interim guild classification for woodland
387 and grassland birds in Australia. *Birds Australia Supplementary Report to State of the Environment*
388 *Report 2006*, Carlton.

389 Thomson, J.R., Moilanen, A.J., Veski, P.A., Bennett, A.F., Nally, R.M., 2009. Where and when to
390 revegetate: a quantitative method for scheduling landscape reconstruction. *Ecological Applications*
391 19, 817-828.

392 Van Teeffelen, A.J.A., Vos, C.C., Opdam, P., 2012. Species in a dynamic world: Consequences of
393 habitat network dynamics on conservation planning. *Biological Conservation* 153, 239-253.

394 Veríssimo, D., Fraser, I., Groombridge, J., Bristol, R., MacMillan, D.C., 2009. Birds as tourism
395 flagship species: a case study of tropical islands. *Animal Conservation* 12, 549-558.

396 Watson, J., Watson, A., Paull, D., Freudenberger, D., 2003. Woodland fragmentation is causing the
 397 decline of species and functional groups of birds in southeastern Australia. *Pacific Conservation*
 398 *Biology* 8, 261-270.

399 Westgate, M.J., Barton, P.S., Lane, P.W., Lindenmayer, D.B., 2014. Global meta-analysis reveals low
 400 consistency of biodiversity congruence relationships. *Nat Commun* 5.

401 Wiens, J.A., Hayward, G.D., Holthausen, R.S., Wisdom, M.J., 2008. Using Surrogate Species and
 402 Groups for Conservation Planning and Management. *Bioscience* 58, 241-252.

403 Williams, P.H., Araújo, M.B., 2000. Using probability of persistence to identify important areas for
 404 biodiversity conservation. *Proceedings of the Royal Society of London. Series B: Biological Sciences*
 405 267, 1959-1966.

406 Williams, P.H., Gaston, K.J., 1994. Measuring more of biodiversity: Can higher-taxon richness
 407 predict wholesale species richness? *Biological Conservation* 67, 211-217.

408 Yong, D.L., Barton, P.S., Okada, S., Crane, M., Lindenmayer, D.B., 2016. Birds as surrogates for
 409 mammals and reptiles: Are patterns of cross-taxonomic associations stable over time in a human-
 410 modified landscape? *Ecological Indicators* 69, 152-164.

411

Table 1. Total richness, total patches occupied, and median and range of patches occupied by each taxonomic group, and the area of the best patch sets selected to meet the 25%, 50% and 75% representation targets of species occurrences.

	Taxon		Species		Area (ha) of best patch set		
	Total richness	Total patches occupied	Median patches occupied	Range patches occupied	25% target	50% target	75% target
Woodland birds	72	189	10.25	1 - 157	274.10	451.10	917.16
Listed birds	10	106	4.25	1 - 43	80.90	187.80	448.76
Arboreal marsupials	3	96	38.00	2 - 59	54.30	168.00	420.96
Reptiles	12	168	5.00	1 - 90	205.40	352.06	719.36




Fig. 1. Spatial dissimilarity of best patch sets selected for 25%, 50% and 75% representation targets of all woodland birds, listed woodland birds, arboreal marsupials and reptiles.

■ Woodland birds ▲ Listed birds ◆ Arboreal marsupials ● Reptiles




Fig. 2. Minimum occurrence for each taxon achieved by the best patch sets for: (a) woodland birds, (b) listed woodland birds, (c) arboreal marsupials, and (d) reptiles. Points represent 25%, 50% and 75% representation targets. Point color indicates if representation target was met (black) or unmet (grey).

Supplementary material

Effectiveness of woodland birds as taxonomic surrogates in conservation planning for biodiversity on farms

Karen Ikin*, Ding Li Yong and David Lindenmayer¹

Email: karen.ikin@anu.edu.au

Table A1. Average (minimum – maximum) site attributes in remnant woodland, regrowth woodland, and planting patches investigated in this study.

Attribute	Remnant woodland	Regrowth woodland	Planting
Patch area (ha)	7.8 (0.8 - 44.7)	10.57 (0.5 - 53.8)	4.24 (0.3 - 60.3)
Canopy height (m)	20.64 (7.67 - 30)	18.4 (8 - 32.5)	10.33 (0.33 - 21.67)
Number of trees	6.3 (0.67 - 52.67)	14.83 (1.33 - 58.33)	29.85 (0 - 203.67)
Number of trees >0.5 m DBH	18.18 (0 - 45.76)	10.42 (0 - 32.24)	1.19 (0 - 11.44)
Number of dead trees/ha	14.49 (0 - 75)	39.89 (0 - 291.67)	17.08 (0 - 100)
Number of strata	2.18 (1.33 - 3.67)	2.5 (1.67 - 4)	2.48 (1.33 - 3.67)
Number of mistletoe/ha	8.33 (0 - 166.67)	8.06 (0 - 158.33)	3.28 (0 - 66.67)
Log density (m ³ /ha)	198.91 (0 - 1100)	168.99 (0 - 766.67)	32.65 (0 - 283.33)
% Overstory cover	30.87 (0 - 65)	30.04 (6.67 - 80)	14.96 (0 - 86.67)
% Midstory cover	0.24 (0 - 11.67)	0.66 (0 - 10)	7.02 (0 - 71.67)
% Understorey cover	1.11 (0 - 18.33)	1.61 (0 - 15)	1.98 (0 - 11.67)
% Rock cover	4.15 (0 - 41.67)	4.45 (0 - 33.33)	1.24 (0 - 13.33)
% Native tussock cover	11.02 (0 - 62.08)	11.28 (0 - 38.75)	4.49 (0 - 35.42)
% Annual grasses cover	32.49 (0 - 176.25)	25.85 (0 - 74.58)	25.23 (0 - 90.83)
% Broad leaf weeds cover	2.54 (0 - 26.67)	2.42 (0 - 20)	2.38 (0 - 22.08)
% Forbs cover	5.45 (0 - 23.75)	4.52 (0 - 21.25)	3.81 (0 - 30.83)
% Moss and lichen cover	3.25 (0 - 25.67)	3.28 (0 - 24.17)	0.86 (0 - 9.17)
% Bare earth	16.86 (0 - 59.58)	14.06 (0 - 46.67)	16.67 (0 - 68.58)
% Leaf litter	27.56 (0.42 - 69.17)	35.18 (2.08 - 77.5)	37.76 (0.83 - 89.25)

Table A2. List of species in each taxon included in the analyses and the number of patches occupied in each year. Taxonomy follows Christidis & Boles (2008) for woodland birds, Jackson & Groves (2015) for arboreal marsupials, and Wilson & Swan (2013) for reptiles.

Taxon	Scientific name	Common name	# Patches	
			2008	2011
Woodland birds (<u>Listed birds</u>)				
	<i>Acanthiza chrysorrhoa</i>	Yellow-rumped thornbill	38	40
	<i>Acanthiza lineata</i>	Striated thornbill	3	8
	<i>Acanthiza nana</i>	Yellow thornbill	14	27
	<i>Acanthiza reguloides</i>	Buff-rumped thornbill	10	12
	<i>Anthochaera carunculata</i>	Red wattlebird	61	41
	<i>Aphelocephala leucopsis</i>	Southern whiteface	5	0
	<i>Artamus cyanopterus</i>	Dusky woodswallow	25	19
	<i>Artamus personatus</i>	Masked woodswallow	10	9
	<i>Cacomantis pallidus</i>	Pallid cuckoo	2	0
	<i>Chalcites baslis</i>	Horsfield's bronze-cuckoo	7	6
	<i>Chalcites lucidus</i>	Shining bronze-cuckoo	0	3
	<u><i>Chthonicola sagittata</i></u>	<u>Speckled warbler</u>	1	6
	<u><i>Climacteris picumnus</i></u>	<u>Brown treecreeper</u>	43	28
	<i>Colluricincla harmonica</i>	Grey shrike-thrush	65	54
	<i>Coracina novaehollandiae</i>	Black-faced cuckoo-shrike	57	61
	<i>Corcorax melanorhamphos</i>	White-winged chough	49	49
	<i>Cormobates leucophaea</i>	White-throated treecreeper	9	9
	<i>Cracticus nigrogularis</i>	Pied butcherbird	24	18
	<i>Cracticus tibicen</i>	Australian magpie	157	141
	<i>Cracticus torquatus</i>	Grey butcherbird	18	19
	<i>Dacelo novaeguineae</i>	Laughing kookaburra	27	28
	<i>Dicaeum hirundinaceum</i>	Mistletoebird	11	12
	<i>Entomyzon cyanotis</i>	Blue-faced honeyeater	11	6
	<i>Eopsaltria australis</i>	Eastern yellow robin	1	1
	<i>Eurystomus orientalis</i>	Dollarbird	2	1
	<i>Falcunculus frontatus</i>	Crested shrike-tit	29	25
	<i>Geopelia placida</i>	Peaceful dove	21	7
	<i>Gerygone albogularis</i>	White-throated gerygone	2	7
	<i>Gerygone fusca</i>	Western gerygone	5	12
	<u><i>Glossopsitta pusilla</i></u>	<u>Little lorikeet</u>	8	0
	<i>Lalage sueurii</i>	White-winged triller	32	55
	<i>Lichenostomus chrysops</i>	Yellow-faced honeyeater	3	6
	<i>Lichenostomus fuscus</i>	Fuscous honeyeater	5	6
	<i>Lichenostomus penicillatus</i>	White-plumed honeyeater	115	103
	<i>Malurus cyaneus</i>	Superb fairy-wren	41	76
	<i>Manorina melanocephala</i>	Noisy miner	106	82
	<u><i>Melanodryas cucullata</i></u>	<u>Hooded robin</u>	5	0
	<i>Melithreptus brevirostris</i>	Brown-headed honeyeater	6	9

Taxon	Scientific name	Common name	# Patches	
			2008	2011
	<i>Melithreptus gularis</i>	<u>Black-chinned honeyeater</u>	3	6
	<i>Melithreptus lunatus</i>	White-naped honeyeater	0	2
	<i>Microeca fascinans</i>	Jacky winter	17	13
	<i>Myiagra inquieta</i>	Restless flycatcher	9	18
	<i>Myiagra rubecula</i>	Leaden flycatcher	5	3
	<i>Neochmia temporalis</i>	Red-browed finch	2	2
	<i>Neophema pulchella</i>	<u>Turquoise parrot</u>	0	2
	<i>Ocyphaps lophotes</i>	Crested pigeon	92	53
	<i>Oriolus sagittatus</i>	Olive-backed oriole	3	4
	<i>Pachycephala rufiventris</i>	Rufous whistler	22	57
	<i>Pardalotus punctatus</i>	Spotted pardalote	4	10
	<i>Pardalotus striatus</i>	Striated pardalote	109	97
	<i>Petroica boodang</i>	<u>Scarlet robin</u>	1	1
	<i>Petroica goodenovii</i>	Red-capped robin	11	7
	<i>Phaps chalcoptera</i>	Common bronzewing	19	8
	<i>Philemon citreogularis</i>	Little friarbird	23	12
	<i>Philemon corniculatus</i>	Noisy friarbird	6	10
	<i>Platycercus elegans</i>	Crimson rosella	24	12
	<i>Platycercus eximius</i>	Eastern rosella	147	126
	<i>Polytelis swainsonii</i>	<u>Superb parrot</u>	37	20
	<i>Pomatostomus superciliosus</i>	White-browed babbler	10	4
	<i>Pomatostomus temporalis</i>	<u>Grey-crowned babbler</u>	6	5
	<i>Psephotus haematonotus</i>	Red-rumped parrot	110	85
	<i>Rhipidura albiscapa</i>	Grey fantail	7	31
	<i>Rhipidura leucophrys</i>	Willie wagtail	126	110
	<i>Sericornis frontalis</i>	White-browed scrubwren	3	3
	<i>Smicrornis brevirostris</i>	Weebill	21	40
	<i>Stagonopleura guttata</i>	<u>Diamond firetail</u>	9	19
	<i>Strepera graculina</i>	Pied currawong	5	3
	<i>Struthidea cinerea</i>	Apostlebird	1	1
	<i>Taeniopygia bichenovii</i>	Double-barred finch	1	1
	<i>Taeniopygia guttata</i>	Zebra finch	1	1
	<i>Todiramphus sanctus</i>	Sacred kingfisher	22	16
	<i>Zosterops lateralis</i>	Silvereye	5	9
Arboreal marsupials				
	<i>Petaurus norfolcensis</i>	Squirrel glider	2	7
	<i>Pseudocheirus peregrinus</i>	Common ringtail possum	37	39
	<i>Trichosurus vulpecula</i>	Common brushtail possum	37	59
Reptiles				
	<i>Aprasia parapulchella</i>	Pink-tailed worm lizard	1	28
	<i>Carlia tetradactyla</i>	Southern rainbow skink	32	1
	<i>Chelodina longicollis</i>	Long-necked turtle	0	26
	<i>Christinus marmoratus</i>	Southern marbled gecko	23	49
	<i>Cryptoblepharus pannosus</i>	Ragged snake-eyed skink	52	20

Taxon	Scientific name	Common name	# Patches	
			2008	2011
	<i>Ctenotus spaldingi</i>	Spalding's ctenotus	9	4
	<i>Delma inornata</i>	Olive legless lizard	20	8
	<i>Diplodactylus vittatus</i>	Eastern stone gecko	9	14
	<i>Egernia striolata</i>	Tree crevice-skink	14	5
	<i>Hemiergis talbingoensis</i>	Three-toed skink	6	1
	<i>Lampropholis guichenoti</i>	Garden skink	1	1
	<i>Lerista bougainvillii</i>	Bougainville's skink	2	4
	<i>Lerista timida</i>	Three-toed lerista	4	2
	<i>Menetia greyii</i>	Grey's skink	3	0
	<i>Morelia spilota</i> ssp. <i>metcalfei</i>	Inland carpet python	0	90
	<i>Morethia boulengeri</i>	Boulenger's skink	87	15
	<i>Pogona barbata</i>	Eastern bearded dragon	3	1
	<i>Pseudechis porphyriacus</i>	Red-bellied black snake	0	7
	<i>Pseudonaja textilis</i>	Eastern brown snake	3	0
	<i>Tiliqua scincoides</i> ssp. <i>scincoides</i>	Eastern blue-tongue	3	1
	<i>Underwoodisaurus milii</i>	Barking gecko	1	3
	<i>Varanus varius</i>	Lace monitor	2	0

References

- Christidis, L. & Boles, W. (2008). *Systematics and taxonomy of Australian birds*, CSIRO Publishing, Collingwood.
- Jackson, S. & Groves, C. (2015). *Taxonomy of Australian mammals*, CSIRO Publishing, Clayton South.
- Wilson, S. & Swan, G. (2013). *Complete guide to reptiles of Australia (4th edn)*. Reed New Holland, Sydney.




Figure A1. Map of the study area in the South West Slopes, Australia.




Figure A2. Minimum occurrence for each taxon achieved by the best patch sets for: (a) woodland birds including subset of listed birds, and (b) woodland birds excluding subset of listed birds. Points represent 25%, 50% and 75% representation targets. Point color indicates if representation target was met (black) or unmet (grey).